

**Previous Year (2018)
Question Paper of
Numerical Methods
BCA-0602**



**You can also get other previous year
question papers of BCA (HPU) on
www.read4bca.com**

Roll No. ...6115XXZ.W440001

Total No. of Questions : 9]
(1048)

[Total No. of Printed Pages : 8

**B.C.A. (CBCS) RUSA Vith Semester
Examination**

4038

NUMERIAL METHODS

Paper : BCA0602

Time : 3 Hours]

[Maximum Marks : 70

Note :- Attempt *four* questions in all, selecting *one* question from each of the Sections B, C, D and E. Question No. 1 is Section A is compulsory.

Section-A

1. (A) Answer all the following *ten* questions with 1 mark each on the answer-book.

(i) The order of convergence in Newton-Raphson method is :

(a) 2

(b) 3

C-670

(1)

Turn Over

- (c) 0
- (d) None of these
- (ii) Which of the following relation is false ?
- (a) $E = 1 + \Delta$
- (b) $E^{-1} = 1 - \nabla$
- (c) $\nabla^2 = 1 - 2E + E^{-2}$
- (d) None of these
- (iii) In Gauss-elimination method for solving a system of linear algebraic equations, triangularization leads to :
- (a) Diagonal matrix
- (b) Lower triangular matrix
- (c) Upper triangular matrix
- (d) Singular matrix
- (iv) Which of the following methods always converges to root of equation $f(x) = 0$?
- (a) Newton-Raphson method
- (b) Regula-Falsi method

(c) Secant method

(d) None of these

(v) By Simpson's (1/3)- rule, the value of

$$\int_1^7 \frac{1}{x} dx \text{ is :}$$

(a) 1.358

(b) 1.958

(c) 1.625

(d) 1.458

(vi) The Gauss-Seidal method gives results faster when the pivotal elements are :

(a) Smaller than other coefficients

(b) Larger than other coefficients

(c) Equal to other coefficients

(d) None of these

(vii) The value of $\Delta^2 x^3$ at $x = 0$, is

(viii) $(0.735816E4) + (0.635742E4) = \dots\dots\dots$

(ix) $(0.999658E-3) - (0.994576E-3) = \dots\dots\dots$

(x) The first term of the series whose second and subsequent terms are 8, 3, 0, -1, 0 is
..... (1×10=10)

Short answer type questions :

(B) Answer all *four* questions. Each question carries 5 marks.

(i) Evaluate the sum $S = \sqrt{3} + \sqrt{5} + \sqrt{7}$ to four significant digits and find relative error..

(ii) Find the root of the equation $e^4 = 4x$, which is approximately 2, correct to three places of decimals.

(iii) Express $3x^3 - 4x^2 + 3x - 11$, in factorial notation.

(iv) Derive Simpson's 1/3-rule using method of undetermined coefficients. (4×5=20)

Section-B

2. (a) Given $f(x) = \sin x$, construct the Taylor series approximations of order 0 to 7 at $x = \frac{\pi}{3}$ and state their absolute errors. (5)
- (b) If $z = \frac{1}{8}xy^3$, find the percentage error in z when $x = 3.14 \pm 0.0016$ and $y = 4.5 \pm 0.05$. (5)
3. (a) If $p = 3c^6 - 6c^2$, find the percentage error in p at $c = 1$, if the error in c is 0.005. (5)
- (b) Convert the following binary numbers to decimal form :
(100101)₂ and (10000001)₂. (5)

Section-C

4. (a) Find a root of the equation $4e^{-x} \sin x - 1 = 0$ by Regula-Falsi method given that the root lies between 0 and 0.5. (5)

(b) Find the roots of the equation $\sin x = 1 + x^3$, between $\frac{0}{-2}$ and $\frac{0}{-1}$ correct to three decimal place by Newton-Raphson method. (5)

5. Solve the system $6x + y + z = 20$, $3x + 4y - z = 6$, $x - y + 5z = 7$ using both Jacobi and Gauss-Seidel methods. (10)

Section-D

6. (a) Using the method of separation of symbols, show that $\Delta^n u_{x-n} = u_x - nu_{x-1} + \frac{n(n-1)}{2}u_{x-2} + \dots + (-1)^n u_{x-n}$. (5)

(b) Using Newton forward difference formula, find the sum

$$S_n = 1^3 + 2^3 + \dots + n^3. \quad (5)$$

7. (a) From the following table, find the value of $e^{1.17}$ using Gauss's forward formula : (5)

x	1.00	1.05	1.10	1.15	1.20	1.25	1.30
e^x	2.7183	2.8577	3.0042	3.1582	3.3201	3.4903	3.6693

- (b) In an examination the number of candidates who obtained marks between certain limits were as follows :

Mark	30-40	40-50	50-60	60-70	70-80
No. of Students	31	42	51	35	31

Find the number of candidates whose scores lie between 45 and 50. (5)

Section-E

8. (a) From the following values of x and $y = f(x)$:

x	0.4	0.5	0.6	0.7	0.8
$f(x)$	1.5836	1.7974	2.0442	2.3275	2.6511

Find $\frac{dy}{dx}$ at $x = 0.6$. (5)

- (b) The function $y = 3xe^{-x}$ is tabulated below.

(3,0.4481), (4,0.2198), (5,0.1011).

Find $y'(x)$ at $x = 3, 4$ and 5 and compare your results with the exact values. (5)

9. (a) Derive Simpson's 3/8 rule and using this rule

evaluate $\int_0^1 \frac{1}{1+x} dx$ with $h = \frac{1}{6}$. (5)

(b) Compute the integral $\int_0^{\pi/2} \sqrt{1 - 0.162 \sin^2 x} dx$
by Weddle's rule. (5)



You can also get other previous year
question papers of BCA (HPU) on

www.read4bca.com

Thank You

